TECHNICAL ASSISTANCE TEAM FOR EMERGENCY RESPONSE REMOVAL AND PREVENTION EPA CONTRACT 68-01-7367

Mr. Steven J. Faryan
Deputy Project Officer
Emergency Response Section
Western Response Unit
U.S. Environmental Protection Agency
11th Floor
230 South Dearborn Street
Chicago, Illinois 60604

January 15, 1988

TAT-05-G2-00268

EPA Region 5 Records Ctr.

225736

Re: Lusher Street, Elkhart, Indiana

TDD# 5-8710-16

Dear Mr. Faryan:

The U.S. Environmental Protection Agency tasked the Technical Assistance Team on October 31, 1987, to investigate alleged ground water contamination in a mixed residential/industrial area of Elkhart, Indiana. Information gathered prior to the site investigation indicated that the affected Lusher Street area included fifteen water wells which contained trichloroethene and/or 1,1,1 trichloroethane in concentrations potentially threatening to human health. This report includes a review of background information, a discussion of analytical results, and recommendations for possible immediate action.

On November 17, 1987 cost estimates were presented to Deputy Project Officer Steve Faryan. These cost estimates addressed the immediate installation of eleven point-of-use carbon filters, two 3.5 cubic foot carbon filters, and hook-up of two residences to city water.

For the purposes of this Removal Action Plan two phases are presented.

- o Phase I entails providing immediate alternate drinking water supplies to thirteen residences and businesses and hook-up of two severely affected residences to city water. This phase could be completed at a total cost of \$15,674.25 (ERCS).
- o Phase II entails conducting an extent-of-contamination study and possible hook-up of fifty additional residences and businesses to city water. This phase could be completed at a cost of \$85,290.00 (ERCS).

Roy F. Weston, Inc.
SPILL PREVENTION & EMERGENCY RESPONSE DIVISION

In Association with ICF Technology Inc., C.C. Johnson & Malhotra, P.C., Resource Applications, Inc., Geo/Resource Consultants, Inc., and Environmental Toxicology International, Inc.



Mr. Steven J. Faryan

-2-

January 15, 1988

As such, the entire corrective action will cost an estimated \$130,000.

Should you have any questions or require additional information, please feel free to contact us.

Very truly yours,

ROY F. WESTON, INC.

Jeffrey S. Binkley

Environmental Geologist

Fig Scott D. Springer

Technical Assistance Team

Leader, Region V

JSB/ljs

Attachment

18.

X

Luder St. Filo 01 0003

LUSHER STREET

GROUND WATER INVESTIGATION

ELKHART, INDIANA

Prepared For:

U.S. Environmental Protection Agency
Region V
230 South Dearborn Street
Chicago, Illinois

CONTRACT NO. 68-01-7367

TAT-05-N-00045

TDD NO. 5-8710-16

Prepared By:

WESTON-SPER
Technical Assistance Team
Region V

January 1988

TABLE OF CONTENTS

		<u>Page</u>
List	of Figures	.iii
List	of Tables	iv
1.0	Site Description	1
2.0	Site Background	1
3.0	Site Assessment	4
4.0	Analytical Results	4
5.0	Threats to Human Health and the Environment	8
6.0	Recommendations	9
7.0	Total Project Cost Summary	12

. **.**

LIST OF FIGURES

					<u>Pa</u>	ge
Figure	1	-	Site	Location	Map	. 2
Figure	2	_	Site	Мар	• • • • • • • • • • • • • • • • • • • •	. 3

LIST OF TABLES

							<u> </u>	<u>ag</u>	<u>e</u>
Table	1	-	Analytical	Results	of	ECHD	Sampling	• •	5
Table	2	_	Analytical	Results	of	TAT	Sampling	• •	7

1.0 SITE DESCRIPTION


The City of Elkhart is located in Elkhart County in north-central Indiana (Figure 1). The Lusher Street site under investigation encompasses a 46 square block residential-industrial area on the southwest side of Elkhart (Figure 2). The area is bordered to the north by the St. Joseph River, to the west by State Road 19, to the east by Avalon Street (north of Lusher Street) and 15th Street (south of Lusher Street), and to the south by Leininger Street.


The topography of the area is generally flat (0 to 2 percent slopes) with run-off flowing north towards the St. Joseph River. Soils in the area are of the Oshtemo-Fox Association (USDA 1974). In a representative profile the association varies from loamy sand to sandy loam to coarse sand and gravel at a depth of 4 to 5 feet. Oshtemo soils typially have a low available moisture capacity and a low organic matter content. In addition, they are characterized by a moderately rapid permeability and slow run-off. Underlying strata consist of glacial outwash deposits whose average depth is 175 feet. Interbedded within these deposits is a layer of silt and clay whose maximum thickness is 80 feet and average thickness is 20 feet. Where present, this layer divides the outwash deposits into two aquifers and confines the deeper aquifer (Imbrigiotta and Martin, 1981). In the area currently under investigation the confining silt and clay layer is absent. The direction of ground water flow in the area is northeasterly, also towards the St. Joseph River.

The predominant contaminants detected in water wells in this area were trichloroethylene (TCE) (71 to 804 parts per billion (ppb)) and 1,1,1-trichloroethane (1,1,1-TCA) (257 to 3,800 ppb).

2.0 SITE BACKGROUND

While conducting a Potential Responsible Party (PRP) extent-of-contamination survey, Gemeinhardt, Inc., detected contamination in several wells in the four square block area immediately south of Lusher Street. Gemeinhardt, Inc., situated approximately 3/4 mile south of Lusher Street believed the contamination in this area was independent of the plume they were addressing, therefore, no further actions were taken and the Elkhart County Health Department (ECHD) was notified. Subsequently, the ECHD executed an all-inclusive sampling effort within the area bordered by State Road 19 on the west, 15th Street on the east, Leininger Street to the south, and Lusher Street to the north. As preliminary analytical results indicating widespread contamination were received, the sampling effort was expanded to include an area north of Lusher Street. This sampling effort included





all wells within an area bordered by State Road 19 to the west, Avalon Street to the east, Lusher Street to the south and the St. Joseph River to the north.

In total, the ECHD sampled 145 wells identifying 103 which were contaminated with 1,1,1-TCA and/or TCE. Fifteen of the 103 affected wells were found to contain levels that exceeded or were within 50% of the U.S. Environmental Protection Agency (U.S. EPA) removal action level for TCE (128 ppb) and/or 1,1,1-TCA (500 ppb) (Table 1). Subsequently, the ECHD requested assistance from the U.S. EPA in providing alternate drinking water supplies to the affected residents and businesses.

3.0 SITE ASSESSMENT

On October 30, 1987 Technical Assistance Team (TAT) member Jeff Binkley met with ECHD officials and reviewed their analytical results to evaluate the extent of the contaminant plume. The TAT constructed a map of the area indicating ECHD results and developed a sampling plan to verify previous results and begin to define the extent-of-contamination. On November 3, 1987 TAT member Jeff Binkley sampled four of the wells previously sampled by the ECHD to confirm their results and evaluate the immediate threat to human health. The samples were analyzed for volatile organics by Aqualab, Bartlett, Illinois, under TAT Analytical Services TDD# 5-8711-L1.

4.0 ANALYTICAL RESULTS

The analytical results confirmed the ECHD findings at all four locations (Table 2). Of greatest concern were the levels of 1,1,1-TCA (1,590 ppb) at 2121 W. Indiana and TCE (804 ppb) at 2205 17th Street which exceeded the U.S. EPA removal action level and the Agency for Toxic Substance and Disease Registry (ATSDR) recommended bathing standards of 500 ppb and 130 ppb, respectively.

Although TCE and 1,1,1-TCA were present in both areas, TAT and ECHD analytical results indicate that the predominant contaminant north of Lusher Street is 1,1,1-TCA, while south of Lusher Street the prevalent contaminant is TCE. These varying concentrations suggest that multiple plumes or sources are affecting the water quality in the two areas.

TABLE 1

ANALYTICAL RESULTS OF ECHD SAMPLING* LUSHER STREET GROUND WATER CONTAMINATION ELKHART, INDIANA

SEPTEMBER 15 - OCTOBER 29, 1987

	Corresponding Location		
Sample Location	(Figure 2)	Contaminant	Concentration
Neitzke Residence	1	1,1,1-TCA	827 ppb
2121 W. Indiana		TCE	45 ppb
Speraw Residence	2	1,1,1-TCA	276 ppb
1401 El Rono		TCE	21 ppb
Hargrove Residence	3	1,1,1-TCA	97 ppb
1601 Avalon		TCE	160 ppb
Roth Residence	4	1,1,1-TCA	266 ppb
2200 W. Indiana		TCE	4 ppb
Frick Residence	5	1,1,1-TCA	719 ppb
2108 W. Indiana		TCE	30 ppb
Powers Residence	6	1,1,1-TCA	330 ppb
1529 Okema		TCE	4 ppb
Wilsey Residence	7	1,1,1-TCA	572 ppb
2124 W. Indiana		TCE	30 ppb
Blocher Residence 2217 W. Indiana	.8	1,1,1-TCA,	257 ppb
Bentley Residence	9	1,1,1-TCA	358 ppb
1429 El Reno		TCE	16 ppb
Interior Access	10	1,1,1-TCA	263 ppb
1620 Waurika		TCE	9 ppb
Homan Lumber	11	1,1,1-TCA	3,800 ppb
1650 Lusher		TCE	608 ppb
Sotebeer Construction	12	1,1,1-TCA	516 ppb
1747 Lusher		TCE	390 ppb

TABLE 1 (Continued)

Sample Location	Corresponding Location (Figure 2)	Contaminant	Concentration
Glass Master	13	1,1,1-TCA	34 ppb
1720 Markle		TCE	93 ppb
McCreary Residence	14	1,1,1-TCA	24 ppb
1741 Fieldhouse		TCE	71 ppb
Miller Residence	15	1,1,1-TCA	34 ppb
2205 17th Street		TCE	1,390 ppb

^{*}Samples Analyzed by ECHD Lab.

TABLE 2

ANALYTICAL RESULTS OF TAT SAMPLING*

LUSHER STREET GROUND WATER CONTAMINATION

ELKHART, INDIANA

NOVEMBER 3, 1987

Sample Location	Corresponding Location (Figure 2)	Contaminant	Concentration
Miller Residence	15	1,1,1-TCA	3.2 ppb
2205 17th Street		TCE	804 ppb
Neitzke Residence	1	1,1,1-TCA	1,590 ppb
2121 W. Indiana		TCE	38.5 ppb
Hargrove Residence	3	1,1,1-TCA	49 ppb
1601 Avalon		TCE	106 ppb
Speraw Residence	2	1,1,1-TCA	140 ppb
1401 El Reno		TCE	20 ppb

^{*}Samples Analyzed by Aqualab, Bartlett, Illinois.

5.0 THREATS TO HUMAN HEALTH AND THE ENVIRONMENT

5.1 Threats as Related to the National Contingency Plan

Paragraph (b)(2), Section 300.65 of the National Contingency Plan (NCP) outlines several conditions that must be considered to warrant a removal action. These conditions, two of which exist at the Lusher Street area, and which will be elaborated on within the following subsections are:

- Actual or potential exposure to hazardous substances or pollutants or contaminants by nearby populations, animals or food chain; and,
- o Actual or potential contamination of drinking water supplies or sensitive ecosystems.

Based on the preceding considerations, contaminated water wells in the Lusher Street area pose a substantial and imminent threat to human health.

5.1.1 Actual or Potential Exposure

The presence of TCE and 1,1,1-TCA contaminated ground water beneath the Lusher Street area has been documented by both the TAT and the ECHD. These contaminants pose a threat of exposure as local residents and businesses utilize ground water from the contaminated aquifer.

5.1.2 <u>Actual or Potential Contamination of Drinking Water Supplies</u>

Ground water contamination has been documented in the Lusher Street area by both the TAT and the ECHD. Of 103 affected wells, 11 were found to contain levels exceeding or within 50% of the U.S. EPA's removal action levels. In addition, 4 wells contained levels exceeding ATSDR recommended bathing standards. Given the local geological conditions and contaminants present in the area, the potential for further contamination of the local ground water reservoir exists.

5.2 Threats as Related to Specific Chemical Contaminants Located On-Site

The TCE and 1,1,1-TCA contamination recently documented in the Lusher Street area poses health threats via direct contact, inhalation, and ingestive routes of exposure.

5.2.1 <u>1,1,1-Trichloroethane</u>

1,1,1-TCA, a non-carcinogenic volatile organic compound, affects the skin, eyes, cardiovascular system and central nervous system. Liquid and vapor 1,1,1-TCA are irritating to eyes on contact and may cause conjunctivitis. Repeated skin contact may produce a dry, scaly, and fissured dermatitis. In addition, 1,1,1-TCA acts as a narcotic and depresses the central nervous system. Acute exposure symptoms include dizziness, incoordination, drowsiness, decreased reaction time and unconsciousness.

5.2.2 Trichloroethene

TCE, a carcinogenic volatile compound, affects the respiratory system, heart, liver, kidneys, skin, and central nervous system. Exposure to TCE vapor may cause irritation of the eyes, nose and throat. Liquid TCE may cause burning and irritation to the eyes and dermatitis as a result of prolonged skin exposure. Acute exposure to TCE depresses the central nervous sytem exhibiting such symptoms as headache, dizziness, vertigo, tremor, nausea and irregular heartbeat. The U.S. EPA Office of Solid Waste and Emergency Response has assigned TCE a 10⁻⁴ cancer risk level of 280 ppb.

6.0 RECOMMENDATIONS

Because of the imminent threats posed to human health by the Lusher Street area the TAT recommends that a two phase action be implemented in a timely manner. Phase I should provide immediate alternate water supplies to the residents and businesses with wells containing contaminant levels which exceed or are within 50% of the U.S. EPA removal level. Phase II should include an extent-of-contamination study and possible implementation of permanent corrective measures.

6.1 Phase I

Two options were considered for providing alternate drinking water supplies, bottled water and point-of-use carbon filters. Cost comparisons and logistics indicated that installation of point-of-use carbon filters was the preferred option at 11 locations. At two locations, where contaminant levels exceed the capabilities of point-of-use carbon filters, the TAT recommends the installation of larger carbon filters. In addition, the TAT recommends that the two residents affected by contaminant levels exceeding ATSDR bathing standards be immediately hooked-up to city water.

6.1.1 Costs for Phase I

Subcontracted

	<u>Items</u>		Amount
	Point-of-Use Carbon Filters 25.00/filter, installed		\$ 2,475.00
	5 Cubic Foot Carbon Filters 50.00/filter, installed		1,500.00
	ty Hook-Ups .,500.00/hook-up		3,000.00
3% C	DDC		209.25
		Sub-Total	\$ 7,184.25

TAT Costs

Redacted-information not relevant to selection of removal action.

∞ .	Sub-Total	\$ 4,850.00
EPA Costs		
Items		Amount
40 hrs. x \$91.00/hr.	•	\$ 3,640.00
	Sub-Total `~	\$ 3,640.00
6.1.2 Phase I - Cost Summary		
<u>Items</u>		Amount
Subcontracted		\$ 7,184.25
TAT Costs		4,850.00
EPA Costs		3,640.00
	Phase I TOTAL	\$ 15,674.25

6.2 Phase II

Phase II should include an extent-of-contamination study to involve the collection of 50 additional samples to be analyzed for volatile organic analyses. Cost estimates were formulated with the assumption that analytical results would dictate the necessity to hook-up 50 additional residences and businesses to city water.

6.2.1 Phase II Costs

Subcontracted

<u>Items</u>		Amount
50 City Hook-Ups @ \$1,500.00/hook-up		\$ 75,000.00
3% ODC		2,250.00
	Sub-Total	\$ 77,250.00

Analytical

<u>Items</u>		<u>Amount</u>
50 Samples for VOA Analysis @ \$300.00/sample		\$ 15,000.00
e \$300.00/sample	Sub-Total	\$ 15,000.00

TAT Costs

Redacted-information not relevant to selection of removal action.

EPA Costs

<u>Items</u>		Amount
40 hrs. x \$91.00/hr.		\$ 3,640.00
	Sub-Total	\$ 3,640.00

6.2.2 Phase II Cost Summary

<u>Items</u>		Amount
Subcontracted		\$ 77,250.00
TAT Costs		4,400.00
EPA Costs		3,640.00
t .	Phase II TOTAL	\$ 85,290.00
7.0 TOTAL PROJECT COST SUMMARY		
Contractor		
Phase I	\$ 7,184.25	
Phase II 15% Contingency	77,250.00 12,665.14	
Sub-Total	~	\$ 97,099.39
Sub 100ul		Ų <i>31</i> ,033.33
TAT Costs		
Phase I 🔧	\$ 4,850.00	
Phase II	4,400.00	•
Sub-Total	\$ 9,250.00	
15% Contingency	15,952.41	
Extramural TOTAL	4	122,301.80
•		5.
EPA Costs	~	<u>4.</u>
Phase I Phase II	\$ 3,640.00	
	3,640.00	
Intramural TOTAL		7,280.00
	PROJECT TOTAL	\$129,581.80
		OR
		\$130,000.00 =======

REFERENCES

Imbrigiotta, T.E., and A. Martin, Jr. "Hydrologic and Chemical Evaluation of the Ground Water Resources of Northwest Elkhart County, Indiana." USGS Water Resources Division Investigation 81-53. 1981.

USDA Soil Conservation Service, "Soil Survey of Elkhart County." 1974.

48